arXiv:nucl-th/9411027v1 23 Nov 1994

IMPORTANCE OF BARYON-BARYON COUPLING IN HYPERNUCLEI

B. F. Gibson
Theoretical Division, Los Alamos National Laboratory
Los Alamos, New Mexico 87545, U. S. A.
and
Institut für Kernphysik, Forschungszentrum
52425 Jülich, Germany

I. R. Afnan
School of Physical Sciences, The Flinders University of South Australia
Adelaide, South Australia 5001, Australia

J. A. Carlson
Theoretical Division, Los Alamos National Laboratory
Los Alamos, New Mexico 87545, U. S. A.

D. R. Lehman
Department of Physics, The George Washington University
Washington, District of Columbia 20052, U. S. A.

ABSTRACT

The $\Lambda N - \Sigma N$ coupling in Λ–hypernuclei and $\Lambda\Lambda - \Xi N$ coupling in $\Lambda\Lambda$–hypernuclei produce novel physics not observed in the conventional, non-strange sector. Effects of $\Lambda \leftrightarrow \Sigma$ conversion in ${}^{3}_{\Lambda}\mathrm{H}$ are reviewed. The role of $\Lambda N - \Sigma N$ coupling suppression in the $A = 4, 5$ Λ–hypernuclei due to Pauli blocking is highlighted, and the implications for the structure of ${}^{10}_{\Lambda}\mathrm{B}$ are explored. Suppression of $\Lambda\Lambda - \Xi N$ conversion in ${}^{\;\;6}_{\Lambda\Lambda}\mathrm{He}$ is hypothesized as the reason that the $< V_{\Lambda\Lambda} >$ matrix element is small. Measurement of ${}^{\;\;4}_{\Lambda\Lambda}\mathrm{H}$ is proposed to investigate the full $\Lambda\Lambda - \Xi N$ interaction. The implication for $\Lambda\Lambda$ analog states is discussed.

1. Introduction

Since the discovery of the first hyperfragment in a balloon flown emulsion stack some forty years ago, physicists have labored to understand how the addition of the strangeness degree of freedom ($S = -1$ for Λ and Σ and $S = -2$ for Ξ and $\Lambda\Lambda$) alters our picture of nuclei and the baryon-baryon force. Because the Λ and Σ masses differ markedly from the nucleon mass, $SU(3)$ symmetry is broken. How it is broken is a question of importance to our fundamental understanding of the baryon-baryon interaction. Furthermore, the inclusion of an $S \neq 0$ degree of freedom (flavor) in the nucleus adds a third dimension to our evolving picture of nuclei. We have discovered that the physics of hypernuclei is unique — different from the conventional nuclear physics that we have investigated for more than sixty years. Unlike the NN interaction which is dominated by one pion exchange (OPE), the ΛN interaction has no OPE component. Thus, the shorter range aspects of the nuclear force may no longer be hidden under the long range OPE mechanism. $\Lambda N - \Sigma N$ coupling plays a significant role in the hyperon-nucleon (YN) force in contrast to the very limited role that $NN - N\Delta$ coupling appears to play in the $S = 0$ sector. This is due in part to the fact that $m_\Sigma - m_\Lambda \simeq 80$ MeV, whereas $m_\Delta - m_N \simeq 300$ MeV; furthermore, the Δ is a broad resonance that decays via the strong interaction, so that its influence may be spread over a broad energy range. However, it is also possible that baryon coupling within the octet differs from the coupling between members of the baryon octet and the isobar decuplet. Extending the analysis to the $S = -2$ sector of the octet, one would anticipate strong $\Lambda\Lambda - \Xi N$ (YY) coupling; $m_{\Xi N} - m_{\Lambda\Lambda} \simeq 25$ MeV. Pauli blocking of the nucleon in the coupled-channel interaction may well explain why one extracts a relatively weak matrix element from the three observed $\Lambda\Lambda$–hypernuclei, in stark contrast to the strong attraction that one expects because it falls in the same multiplet as the nn interaction.

We will review here some of the interesting aspects of hypernuclei which demonstrate that $S \neq 0$ physics is different. In addition, we will explore how few-baryon systems can be used to constrain our models of the YN and YY interactions, and we will reflect upon what we have so far achieved in understanding the baryon-baryon force in this sector.

2. The Role of $\Lambda N - \Sigma N$ Coupling

We have only to examine the few-body data to see that $\Lambda N - \Sigma N$ coupling plays a significant role in Λ-hypernuclei. In Table 1 we summarize the Λ separation energies,

$$B_\Lambda({}_\Lambda A) = B({}_\Lambda A) - B(A-1) \ ,$$

for the s-shell Λ-hypernuclei [1, 2, 3] along with γ de-excitation energies [4, 5], where they have been measured.

Table 1. Hypernuclear Λ-separation energies and excitation energies in MeV

hypernucleus	B_Λ	E_γ
${}^{3}_{\Lambda}$H	0.13±.05	
${}^{4}_{\Lambda}$H	2.04±.04	1.04±.04
${}^{4}_{\Lambda}$He	2.39±.03	1.15±.04
${}^{5}_{\Lambda}$He	3.10±.02	

The charge symmetry breaking (CSB) in the $A = 4$ isodoublet is obvious: ${}^{4}_{\Lambda}$He is more bound than ${}^{4}_{\Lambda}$H by $\Delta B_\Lambda = 0.35$ MeV. The Λp interaction is stronger than the Λn interaction. [This CSB is almost 3 times that estimated for the ^{3}H – ^{3}He system once the Coulomb energy is taken into account: $B({}^3\mathrm{H}) - B({}^3\mathrm{He}) - E_C \simeq 8.48 - 7.72 - 0.64 \simeq 0.12$ MeV]. One needs no theoretical calculation to see the CSB in the Λ-hypernuclei separation energies, because the Coulomb energy of the ^{3}He core essentially cancels that of ${}^{4}_{\Lambda}$He. The second order Coulomb effect due to the compression of the ^{3}He core in ${}^{4}_{\Lambda}$He by the addition of the Λ actually increases ΔB_Λ by $0.01 - 0.02$ MeV. This CSB provides a clear signal for the importance of $\Lambda N - \Sigma N$ coupling in the YN interaction. The $(\Sigma^+, \Sigma^0, \Sigma^-)$ mass splitting is some 9 MeV, or about 10% of the $\Lambda - \Sigma$ mass difference. Including this alone accounts for a large part of the CSB in the YN interaction.

That $\Lambda N - \Sigma N$ coupling is more significant in Λ-hypernuclei than is $NN - N\Delta$ coupling in conventional nuclei can be seen by comparing Λ separation energies from ${}^{3}_{\Lambda}$H, ${}^{4}_{\Lambda}$He, and ${}^{5}_{\Lambda}$He with neutron separation energies from ^{2}H, ^{3}H, and ^{4}He. In the non-strange sector we know that the ratio of neutron separation energies for neighboring s-shell nuclei is close to a constant:

$$B_n({}^3\mathrm{H})/B_n({}^2\mathrm{H}) \simeq \frac{6}{2} = 3$$

and

$$B_n({}^4\mathrm{He})/B_n({}^3\mathrm{He}) \simeq \frac{20}{6} \simeq 3\,.$$

Thus, one might anticipate that

$$\begin{aligned} B_\Lambda({}^{5}_{\Lambda}\mathrm{He}) &\simeq 3\ B_\Lambda({}^{4}_{\Lambda}\mathrm{He}) \\ &\simeq 6\ \mathrm{MeV} \end{aligned}$$

for model calculations in which $V_{\Lambda N}$ is fitted to the low energy ΛN scattering parameters and reproduces the observed value of $B_\Lambda({}^4_\Lambda\mathrm{He}) \simeq 2$ MeV. This is, in fact, confirmed by several model calculations [6, 7, 8, 9] using central potentials to represent the ΛN force. In contrast, the experimental value for $B_\Lambda({}^5_\Lambda\mathrm{He})$ is 3.1 MeV [1, 2, 3]. Similarly, for the hypertriton, one might estimate

$$\begin{aligned} B_\Lambda({}^3_\Lambda\mathrm{H}) &\simeq 1/3\; B_\Lambda({}^4_\Lambda\mathrm{He}) \\ &\simeq 0.7\,\mathrm{MeV}\,. \end{aligned}$$

This is again confirmed by several model calculations using central potentials to represent the ΛN interaction [10, 11, 12]. In contrast, the experimental value for $B_\Lambda({}^3_\Lambda\mathrm{H})$ is 0.13 MeV.

In other words, our carefully developed intuition coming from the study of conventional, nonstrange nuclei fails to provide a sound means of extrapolating into the strangeness sector. The physics of Λ–hypernuclei is new and different. A large part of this difference can be traced to $\Lambda N - \Sigma N$ coupling. (If $NN - N\Delta$ coupling played as significant a role in the nonstrange sector, then our modeling of nuclei would have required more sophistication. Our shell model calculations, which assume a mean field interaction independent of the quantum numbers of the core, would have required serious modification.) Why do we suggest that a major factor in the anomalously small binding of ${}^5_\Lambda$He is the strong suppression of $\Lambda N - \Sigma N$ coupling? Converting a T=0 Λ into a T=1 Σ requires the simultaneous conversion of the ^{4}He core of ${}^5_\Lambda$He into a T=1, even-parity ${}^4\mathrm{He}^*$ excited state [13]. Such excited states lie high in the continuum. They also have wave functions with more structure than the ^{4}He ground state, which reduces the transition matrix element. Thus, the strength of the full coupled-channel

$$\begin{pmatrix} V_{\Lambda N} & V_{XN} \\ V_{XN} & V_{\Sigma N} \end{pmatrix}$$

interaction is expected to be significantly reduced in ${}^5_\Lambda$He, as the off-diagonal transitions are partially Pauli blocked. That $\Lambda N - \Sigma N$ coupling is suppressed in ${}^5_\Lambda$He is confirmed by Monte Carlo variational calculations [14] using the contemporary Nijmegen soft core potential [15], although the $\Lambda N - \Sigma N$ coupling is so strong in this Nijmegen model that the $A = 5$ system is actually unbound in the calculation. (The anomalously small binding has also been interpreted as a repulsive ΛNN three-body force in models which neglect explicit $\Lambda N - \Sigma N$ coupling [16].)

Similarly, using a YN interaction that incorporates $\Lambda N - \Sigma N$ coupling leads to a reduction in the binding of ${}^3_\Lambda$H compared to that from a single-channel ΛN potential [12]. The hypertriton can be thought of principally as a Λ bound tenuously to a deuteron. The Σ isospin ($T = 1$) differs from that of the Λ ($T = 0$). Therefore, converting the Λ into a Σ requires simultaneously converting the deuteron core into a ${}^1\mathrm{S}_0$ NN pair. The NN ${}^1\mathrm{S}_0$ interaction is much weaker than the tensor dominated ${}^3\mathrm{S}_1$–${}^3\mathrm{D}_1$ force that binds the deuteron. Thus, incorporating $\Lambda N - \Sigma N$ coupling into the interaction model leads to a reduction in the ${}^3_\Lambda\mathrm{H}$ binding. Such an effect might be interpreted as a repulsive ΛNN three-body potential. However, if one formally eliminates the ΣN channel (producing an energy-dependent ΛN potential plus a true ΛNN potential), one finds in explicit calculations that the true ΛNN three-body force is attractive [12]. Nonetheless, it is the case that the anomalously small binding of the hypertriton is the result of strong $\Lambda N - \Sigma N$ coupling in the $S = -1$ baryon-baryon force.

3. The YN Scattering Data

The physics of the $S = -1$ sector clearly differs from that of the $S = 0$ sector. Can we understand the Λ separation energies in terms of the $S = -1$ hyperon–nucleon interaction? Unfortunately the ΛN, ΣN data are very limited. There are some 600 events[17, 18, 19] below 300 MeV/c and another 250 events[20] between 300 and 1500 MeV/c — total cross sections and some angular data. The lack of a significant data base severely inhibits our ability to provide a definitive analysis of hypernuclear data.

The Nijmegen group has taken the lead in modeling [21, 22, 15] the YN potential by fitting a one-boson-exchange (OBE) hypothesis to the combined NN and YN data base using $SU(3)$ constraints. Their pioneering effort to construct such potentials has been summarized by deSwart in a recent Seminar on the hyperon-nucleon interaction [23]. Timmermans elaborated further upon the model [24]. The Jülich group have also produced YN potential models [25, 26, 27], both energy dependent and of the OBE form (in q-space). Reuber discussed the Jülich approach at the same meeting [28]. The models are not the same. For example the F/D ratios differ significantly, which affects the strength in the ΣN channel.

In Table 2 we summarize for several of the models their charge-independent low-energy scattering parameters — the scattering lengths and effective ranges. A quick comparison illustrates the need for better YN data to constrain the potential parameters. Nonetheless, based upon the models, the following observations can be made. The $\Lambda N - \Sigma N$ coupling is strong. The $\Lambda(T = 0) \times N(T = \frac{1}{2})$ direct interaction contains no OPE, although a second order OPE force is generated through $\Lambda N - \Sigma N$ coupling. The short range aspects of the baryon-baryon force, such as heavy meson exchange, play a more important role. Oka recently reviewed the Tokyo quark-cluster modeling of the short range properties of the YN interaction [29]. It is in the YN sector that one begins to find differences between the OBE and quark cluster models of the baryon-baryon force. For example, the OBE potential exhibits strong short range attraction in some ΣN channels, whereas the quark cluster approach appears

Table 2. The charge-independent scattering lengths and effective ranges in fm for the YN potential models listed

Model	Ref.	a^s	r_o^s	a^t	r_o^t
Nijmegen D	[21]	-1.90	3.72	-1.96	3.24
Nijmegen F	[22]	-2.29	3.17	-1.88	3.36
Nijmegen SC	[15]	-2.78	2.88	-1.41	3.11
Jülich A	[27]	-1.56	1.43	-1.59	3.16
Jülich Ã	[27]	-2.04	0.64	-1.33	3.91

to yield consistently short range repulsion.

4. The $A = 4$ Isodoublet

4.1. Charge Symmetry Breaking

We have remarked already that the large CSB in the $A = 4$ isodoublet ground-state binding energies provides strong evidence for the important role of $\Lambda N - \Sigma N$ coupling in the YN interaction. How well can this CSB be understood in terms of the free ΛN interaction? The Nijmegen models predict a noticeable charge symmetry breaking: $V_{\Lambda p} \neq V_{\Lambda n}$. [Recall that strong $\Lambda - \Sigma$ conversion leads to CSB because of the mass difference in the Σ isospin triplet.] The Λp and Λn scattering lengths and effective ranges from Nijmegen model D are compiled in Table 3.

Table 3. The charge-dependent scattering lengths and effective ranges in fm for the Nijmegen model D

channel	a^s	r_0^s	a^t	r_0^t
Λp	-1.77	3.78	-2.06	3.18
Λn	-2.03	3.66	-1.84	3.32

One can use a folding prescription to generate an effective two-body (Λ-nucleus) potential with which to calculate $B_\Lambda({}^4_\Lambda\mathrm{He})$ and $B_\Lambda({}^4_\Lambda\mathrm{H})$. The resulting ΔB_Λ from this approach (in terms of two-body dynamics) using s-wave separable potentials fitted to the low-energy scattering parameters in Table 3 is 0.21 MeV. If one allows for compression of the nuclear core (due to the added binding of the Λ), one can increase this to 0.24 MeV for a 5% core compression. In contrast, using exact four-body equations and the identical separable potentials, one finds a value of $\Delta B_\Lambda = 0.43$ MeV [30]. That is, true 4-body dynamics yields a value for ΔB_Λ about twice that which comes from a model utilizing the same potentials but approximate two-body dynamics.

This perhaps surprising result can be simply understood. One can demonstrate [31, 32] for simple potentials that $|a| > |a'|$ implies that potential V is stronger than potential V' in an n-body calculation, where n = 2, 3, 4. That is, the potential V is uniformly stronger than V', as intuition would lead one to expect because a represents the overall strength of the interaction. In contrast, for $r > r'$, the same relation holds only for $n = 2$. (The larger the potential range, the stronger is the potential in a two-body sense.) However, if $n = 3, 4$, then $r > r'$ means that V is weaker than V'. Recall Thomas' result [33] from 1935: If in a three-body system the range of the force → 0 for one pair, then the three-body binding becomes infinite. Therefore, the CSB in the $A = 4$ isodoublet can be understood in terms of the scattering length and effective range differences inferred from Table 3. The spin-singlet interactions tend to average in the two systems and contribute little to the Λ separation energy difference; on the other hand, the spin-triplet interaction in ${}^4_\Lambda\mathrm{He}$ is pure Λp while that in ${}^4_\Lambda\mathrm{H}$ is pure Λn. We see that Δa and Δr for the spin-triplet potentials will produce compensating effects in a two-body, folding potential approximation (reducing the estimate of CSB in the isodoublet) but will add constructively in a true four-body calculation. Thus, we understand that the difference between the 0.21 MeV and 0.43 MeV quoted above for identical separable potentials comes from the dynamics of the calculations. When one looks at details of few-body systems, exact model calculations can play a crucial role in a correct interpretation.

We note in closing this discussion that tensor forces cannot be neglected; they are known to be less effective in binding few-body systems than are central forces. When a tensor component is included in the the ΛN spin-triplet interaction, the binding energy difference in the $A = 4$ isodoublet is reduced from 0.43 MeV to 0.37 MeV.

4.2. The Excited States

The ${}^4_\Lambda\mathrm{H}/{}^4_\Lambda\mathrm{He}$ system is even more interesting because there exists a particle-stable excited state for each species. The ground states are 0^+, the excited states are 1^+. It is therefore tempting to argue that the measured E_γ transition energies provide a direct determination of the ΛN spin-spin interaction. However, we suggest 1) that is not the case, 2) that $\Lambda N - \Sigma N$ coupling and the complex structure of the $A = 4$ Λ-hypernuclei must be taken into account in any serious calculation, and 3) that the improper modeling of ${}^4_\Lambda$H as $[\Lambda \times {}^3\mathrm{H}]^{[J]}$ is likely the explanation for the "missing" γ-ray in the ${}^{10}_{\Lambda}$B system.

The existence of particle-stable excited states presents a unique opportunity to test our models of the YN interaction. Fitting both the 0^+ and 1^+ states within the same model is a nontrivial task. $\Lambda N - \Sigma N$ coupling, as a crucial feature of the YN interaction, plays a large role. Furthermore, the complex nature of the four-body system is important. The 1^+ state is not just formed from the 0^+ state by a spin flip of the Λ.

Table 4. The ΛN scattering lengths and effective ranges in fm for the Stepien-Rudzka and Wycech model

channel	spin-singlet	spin-triplet
a(fm)	-1.97	-1.95
r_0(fm)	3.90	2.43

Consider a model example in which the Stepien-Rudzka and Wycech [34] separable potential approximation to the Nijmegen model D is employed. The scattering lengths and effective ranges are displayed in Table 4. If one uses effective ΛN potentials fitted to these parameters (*i.e.,* explicit $\Lambda N - \Sigma N$ coupling in the YN interaction is approximated by constructing a ΛN potential that reproduces the scattering length and effective range of the full interaction) in a full four-body calculation, then one obtains for the total binding energies [35]:

$$B(0^+) = 10.7 \text{ MeV}$$

$$B(1^+) = 11.7 \text{ MeV}.$$

The order of the states is inverted; the 1^+ is the ground state. The reason is clear if one looks at the average ΛN potential in the two cases. One finds:

$$0^+ : V_{\Lambda N} = \frac{1}{2} V^S_{\Lambda N} + \frac{1}{2} V^t_{\Lambda N}$$

and

$$1^+ : V_{\Lambda N} = \frac{1}{6} V^S_{\Lambda N} + \frac{5}{6} V^t_{\Lambda N}.$$

Although $V^S_{\Lambda N}$ from Table 4 is stronger than $V^t_{\Lambda N}$ in a two-body sense, it is the effective ranges that control the situation. The scattering lengths are essentially the same. Because $\mathrm{r}_0(^1S_0)$ is significantly larger than $\mathrm{r}_0(^3S_1)$, the potential $V^S_{\Lambda N}$ is weaker than the potential $V^t_{\Lambda N}$ in a full four-body calculation.

However, the nuclear core in ${}^4_\Lambda$H is not an elementary object. It is a ($J = \frac{1}{2}$, $T = \frac{1}{2}$) composite of three nucleons. Converting a $T = 0$ Λ into a $T = 1$ Σ necessarily brings in $T = \frac{3}{2}$ ^{3}H* states. These lie high in the continuum (just as in the case of ^{4}He mentioned previously) *and* have more spatial structure than the ^{3}H ground state. Therefore, the $\Lambda - \Sigma$ conversion in the medium is expected to be suppressed compared to its effect in the free $\Lambda N - \Sigma N$ interaction. Taking an extreme model approximation of dropping the parts of the $V_{\Lambda N-\Sigma N}$ interaction that couple through $T = \frac{3}{2}$ ^{3}H* states, one can construct appropriately modified ΛN effective interactions which yield [35], in the same full four-body calculation:

$$B(0^+) = 9.6 \text{ MeV}$$

$$B(1^+) = 8.2 \text{ MeV}.$$

That is, one obtains a proper ordering of the states. More importantly, this demonstrates that E_γ is not a measure of the ΛN spin-spin interaction.

How realistic are such separable potential calculations? Monte Carlo variational calculations made with the Nijmegen soft core potential have obtained $B(0^+) \simeq$ 1.5 MeV, while the 1^+ state is unbound [14]. The $\Lambda - \Sigma$ conversion suppression is confirmed, although it is too severe to match the data. Our analysis indicates that the YN force modeling problem lies in the sizeable strength in the $V_{\Lambda N-\Sigma N}$ tensor potential in the spin-triplet channel. Second order tensor effects become very important when effectively 2 $\frac{1}{2}$ YN pairs reside in the spin-triplet state. Such effects are well known in few-nucleon physics and play a significant role also in hypernuclei. The $\Lambda N - \Sigma N$ tensor coupling component provides so much of the free interaction strength in the Nijmegen soft core potential model that suppression of that coupling leaves the spin-triplet dominated 1^+ state unbound. An r-space approximation to Jülich $\tilde{\text{A}}$ is needed in order to test that model.

To reiterate, the $0^+ - 1^+$ energy difference does seem to result from complex few-body dynamics and is not a simple measure of the ΛN spin-spin interaction. It is possible to apply this result to understand experiments involving heavier hypernuclei. One can prescribe a scenario based upon the separable potential model results outlined here which may explain the failure to find a $2^- \to 1^-$ γ following the ^{10}B$(K^-, \pi^-){}^{10}_\Lambda$B* reaction [36]. The prediction of such a transition is based upon a shell model calculation in which the $A = 4$ $0^+ - 1^+$ level splitting is used to define the ΛN spin-spin component of the mean field Λ potential [37, 38]. In such a mean field approach, the $0^+ - 1^+$ splitting in $A = 4$ leads to a prediction that the state ordering in ${}^{10}_{\Lambda}$B is such that the ground state is 1^-. The 3^+ g.s. of the target ^{10}B ensures that the 2^- state in ${}^{10}_{\Lambda}$B is produced. Thus, a γ transition from the 2^- excited state to the 1^- ground state should be seen. Chrien *et al.* observed none between 0.1 and 0.9 MeV [36]. However, consider a cluster model in which ${}^{10}_{\Lambda}$B is modeled as an α-α-Λ-p system. The α's are tightly bound. The level structure of ${}^{10}_{\Lambda}$B will likely be determined by the spin dependence of the ΛN interaction, even though the valence proton lies in the p-shell while the Λ lies in the s-shell which reduces the role of their direct spin-spin interaction and brings into play spin-orbit and tensor forces. That is, the state ordering will likely correspond to that of the free YN interaction with no $\Lambda N - \Sigma N$ coupling suppression. (Recall that such an assumption in the $A = 4$ system gave rise to a 1^+ ground state.) Therefore, one would anticipate that the ground state of ${}^{10}_{\Lambda}$B is likely to be 2^- (not 1^-), and no γ should have been seen in the experiment, because the 1^- state is not produced in the reaction. Clearly ${}^{10}_{\Lambda}$B is an important candidate for further investigation, but it does appear that $\Lambda N - \Sigma N$ coupling plays an important role in the structure of Λ–hypernuclei.

5. Hypernuclear Constraints

In addition to the strong constraints that the ${}^{4}_{\Lambda}$H–${}^{4}_{\Lambda}$He isodoublet places upon any model of the YN interaction, ${}^{3}_{\Lambda}$H and ${}^{5}_{\Lambda}$He also constrain various aspects of the YN interaction. The different hypernuclei act as spin filters, emphasizing different components of the YN potential.

5.1. The Hypertriton

Because the YN interaction does not support a two-body bound state [39], as one can see from the scattering lengths in Table 2, the hypertriton in the $S = -1$ sector serves the function of the deuteron. The fact that $B_\Lambda \simeq 130 \pm 50$ keV suggests that the long range properties of this system should dominate. However, the spin-singlet interaction plays the dominant role in the hypertriton

$$V_{\Lambda N} = \frac{3}{4} V^s_{\Lambda N} + \frac{1}{4} V^t_{\Lambda N} \,,$$

so that $B({}^{3}_{\Lambda}H)$ provides a constraint primarily upon that potential component.

$\Lambda N - \Sigma N$ coupling has been known to play an important role in ${}^{3}_{\Lambda}$H since the early calculations of Schick [40]. More recently, we have explored how this translates into a three-body force (3BF) effect when the ΣN channel is formally eliminated [12]. In Fig. 1 are represented three types of diagrams that enter. In (a) we illustrate the usual iteration of the ΛN potential in ${}^{3}_{\Lambda}$H. In (b) we see what is termed the dispersive energy dependence of the ΛN interaction when the ΣN channel is eliminated. It is effectively repulsive, because the energy denominator in the Σ propagator becomes

Fig. 1 Schematic representation of various interactions among the YN constituents of ${}^{3}_{\Lambda}$H: a) a standard contribution to the ΛNN three-body equations, b) an energy-dependent contribution to the ΛN two-body t matrix arising from $\Lambda N - \Sigma N$ coupling, and c) an effective three-body force arising from $\Lambda N - \Sigma N$ coupling.

larger when the kinetic energy of the second nucleon is added:

$$V_{\Lambda N \to \Sigma N} \, \frac{1}{\Delta E} \, V_{\Sigma N \to \Lambda N}.$$

This effect is analogous to the reduction in strength of the NN tensor force when one embeds it in nuclear matter. In (c) we find the true 3BF that results. The 3BF due to $\Lambda N - \Sigma N$ coupling is actually attractive. One is led to ask whether the hypertriton would be bound were it not for this attraction in a realistic potential calculation.

Miyazawa and Glöckle [41] have recently completed a model calculation of ${}^{3}_{\Lambda}$H using the Jülich OBE model $\tilde{\mathrm{A}}$ [27]. They find the system unbound. A 4% increase in the strength of V^{s}_{YN} would produce binding *but* would destroy the model $\tilde{\mathrm{A}}$ description of the YN data. This result is somewhat puzzling — the a, r_o values for the model $\tilde{\mathrm{A}}$ imply significant attraction in a rank-one separable potential ${}^{3}_{\Lambda}$H calculation: $B_\Lambda \simeq 1$ MeV. Why should the short range potential properties play such a significant role in this weakly bound system?

5.2. The ${}^{5}_{\Lambda}$He *Anomaly*

As discussed above, simple estimates for B_Λ would yield values about a factor of 2 larger than the data. Model calculations confirm this [7, 9, 42, 43, 16]. The anomalously small binding would appear to be due to two sources: 1) a tensor force is less effective in binding a few-body system than a central force with the same low energy two-body properties, and 2) $\Lambda N - \Sigma N$ coupling is suppressed because the $T = 1$ Σ must couple to $T = 1$ even parity states high in the α spectrum. Monte Carlo variational calculations for the Nijmegen soft core model support this analysis [14]. These microscopic calculations include explicit Σ degrees of freedom. The NN force was represented by the Nijmegen '78 potential [44] and the Urbana model 7 three-nucleon interaction [45] was included to better represent the ground-state properties of the three- and four-nucleon systems. The YN tensor force was shown to play a significant role; replacing the YN spin-triplet interaction by the central force spin-singlet intercation led to significant overbinding. In contrast to the hypertriton, it is the spin-triplet interaction that dominates

$$V_{\Lambda N} = \frac{1}{4} V^s_{\Lambda N} + \frac{3}{4} V^t_{\Lambda N} \, .$$

Thus, ${}^{5}_{\Lambda}\mathrm{He}$ provides a significant constraint upon that potential component.

6. The $S = -2$ Sector and the ${}^{6}_{\Lambda\Lambda}\mathrm{He}$ Enigma

The doubly-strange hypernuclei of the $S = -2$ sector provide another handle on the importance of coupling to higher-lying channels in the baryon-baryon interaction. The single reported event [46] is controversial [47]. However $B_{\Lambda\Lambda} \simeq 10.6$ yields a value of

$$\begin{aligned} < V_{\Lambda\Lambda} > &= B_{\Lambda\Lambda} - 2B_{\Lambda}({}^{5}_{\Lambda}\mathrm{He}) \\ &\simeq 4.4 \text{ MeV} \, , \end{aligned}$$

which is consistent with the values of $4 - 5$ MeV extracted from the other two $\Lambda\Lambda$ hypernuclei [51, 52, 53]. Furthermore, Hartree-Fock calculations [48] established that comparable $V_{\Lambda\Lambda}$ and $V_{\Lambda N}$ interactions were required to account for the binding energies of ${}^{6}_{\Lambda\Lambda}\mathrm{He}$ and ${}^{4}_{\Lambda}\mathrm{He}$, respectively. That is, one finds

$$< V_{\Lambda\Lambda} >_{A=6} \simeq < V_{\Lambda N} >_{A=4}$$

could account for the binding energy of both systems. The ΛN interaction is only weakly attractive, which implies a similar property for the $\Lambda\Lambda$ force. The preferred values of the scattering length and effective range of the phenomenological $\Lambda\Lambda$ potential required to reproduce the binding energies of the $\Lambda\Lambda$ hypernuclei in the analysis by Bodmer *et al.* [49] compared to those for the ΛN system listed in Table 2 lead to a similar conclusion, as does the G-matrix analysis of Bando [50] based upon the Nijmegen model D.

However, just as $\Lambda N - \Sigma N$ coupling suppression appears to play a significant role in the ${}^{4}_{\Lambda}$He–${}^{4}_{\Lambda}$H level ordering and in the anomalously small binding of ${}^{5}_{\Lambda}$He, we hypothesize that $\Lambda\Lambda - \Xi N$ coupling is likely to be suppressed in ${}^{\;\,6}_{\Lambda\Lambda}$He, so that only $< V_{\Lambda\Lambda} >$ is sampled. The $\Lambda\Lambda - \Xi N$ interaction lies in the same multiplet as the nn interaction, which almost binds. Thus, it is rather surprising that $< V_{\Lambda\Lambda} >$, as extracted from ${}^{\;\,6}_{\Lambda\Lambda}$He, is weak by comparison. Suppression of the ΞN conversion is a prime candidate to explain this observation. One anticipates such suppression in ${}^{\;\,6}_{\Lambda\Lambda}$He because conversion from $\Lambda\Lambda \rightarrow \Xi N$ in a relative ${}^{1}\mathrm{S}_{0}$ state such that the resulting ΞN pair reside in the 1s shell requires excitation of the ^{4}He core. Five nucleons cannot exist in the 1s shell.

Therefore, we suggest that ${}^{\;\,6}_{\Lambda\Lambda}$He probes primarily the diagonal element $V_{\Lambda\Lambda}$ of the $\Lambda\Lambda - \Xi N$ interaction, while it would be ${}^{\;\,4}_{\Lambda\Lambda}$H (or ${}^{\;\,5}_{\Lambda\Lambda}$He) that would probe the full $\Lambda\Lambda - \Xi N$ force. The $\Lambda\Lambda np$ system can easily couple to ΞNNN in a manner that leaves all four baryons in the 1s shell. Because ${}^{3}_{\Lambda}$H is bound, ${}^{\;\,4}_{\Lambda\Lambda}$H will also be bound. If the full $\Lambda\Lambda - \Xi N$ interaction strength is comparable to that of the nn interaction, then one could even envision a 0^{+} particle-stable excited state as well as a 1^{+} ground state for ${}^{\;\,4}_{\Lambda\Lambda}$H.

7. Summary

In brief conclusion, we reiterate the important physics differences between conventional, nonstrange nuclear physics and hypernuclear physics. The few-baryon systems provide filters through which one can look at particular aspects of the YN interaction. They provide specific constraints on models of the interaction, and are particularly sensitive to $\Lambda N - \Sigma N$ coupling and its induced spin dependence. Furthermore, extrapolating our experience in the $S = -1$ sector to the $S = -2$ sector leads us to predict strong suppression of $\Lambda\Lambda - \Xi N$ coupling in all but the lightest of the $\Lambda\Lambda$ hypernuclei.

Acknowledgement

The work of BFG and JAC was performed under the auspices of the U. S. Department of Energy, while that of DRL was supported in part by the U. S. Department of Energy under Grant No. DE-FG05-86ER40270, and that of IRA was supported by the Australian Research Council. BFG gratefully acknowledges a Research Award for Senior Scientists by the Alexander von Humboldt Stiftung which made possible his stay with the I. K. P. Theorie group at the Forschungszentrum Jülich.

References

[1] M. Juric, G. Bohm, J. Klabuhn, U. Krecker, F. Wysotzki, G. Coremans-Bertand, J. Sacton, G. Wilquet, T. Cantrell, F. Esmael, A. Montwill, D. H. Davis, *et al.* Nucl. Phys. **B52**, 1 (1973).

[2] D. H. Davis and J. Sacton, in *High Energy Physics*, Vol. II, (Academic Press, New York, 1967) p. 365.

[3] D. H. Davis, Proceedings of the *LAMPF Workshop on* (π, K) *Physics*, A. I. P. Conf. Proc. **224**, ed. by B. F. Gibson, W. R. Gibbs, and M. B. Johnson, (American Institute of Physics, New York, 1991), pp. 38-48.

[4] A. Bamberger, M. A. Faessler, U. Lynn, H. Piekarz, J. Piekarz, J. Pniewski, B. Povh, H. G. Ritter, and V. Soergel, Nucl. Phys. **B60**, 1 (1973).

[5] M. Bejidian, E. Descroix, J. Grossiord, A. Guichard, M. Gusakow, M. Jacquin, M. Kudla, H. Piekarz, J. Piekarz, J. Pizzi, and J. Pniewski, Phys. Lett. B **83**, 252 (1979).

[6] A. R. Bodmer, Phys. Rev. **141**, 1387 (1966).

[7] R. C. Herndon, Y. C. Tang, Phys. Rev. **153**, 1091 (1967); **159**, 853 (1967); **165**, 1093 (1969); R. H. Dalitz, R. C. Herndon, and Y. C. Tang, Nucl. Phys. **B47**, 109 (1972).

[8] B. F. Gibson, A. Goldberg, and M. S. Weiss, Phys. Rev. C **6**, 741 (1972).

[9] A. Gal, Adv. Nucl. Physics **8**, 1 (1975).

[10] J. Dabrowski and E. Fedorynska, Nucl. Phys. **A210**, 509 (1973).

[11] B. F. Gibson and D. R. Lehman, Phys. Rev. C **11**, 29 (1975).

[12] I. R. Afnan and B. F. Gibson, Phys. Rev. C **40**, R7 (1989); **41**, 2787 (1990).

[13] B. F. Gibson, A. Goldberg, and M. S. Weiss, Phys. Rev. **181**, 1486 (1969).

[14] J. A. Carlson, Proceedings of the *LAMPF Workshop on* (π, K) *Physics*, A. I. P. Conf. Proc. **224**, ed. by B. F. Gibson, W. R. Gibbs, and M. B. Johnson, (American Institute of Physics, New York, 1991), pp. 198 - 210.

[15] P. M. M. Maessen, T. A. Rijken and J. J. de Swart, Phys. Rev. C **40**, 2226 (1989).

[16] A. R. Bodmer and Q. N. Usmani, Nucl. Phys. **A477**, 621 (1988); Phys. Rev. C **31**, 1400 (1985); A. R. Bodmer, Q. N. Usmani, and J. A. Carlson, Phys. Rev. C **29**, 684 (1984).

[17] G. Alexander, U. Karshon, A. Shapira, G. Yakutieli, R. Engelmann, H. Filthuth, A. Fridman, and A. Minguzzi-Ranzi, Phys. Rev. Lett. **13**, 484 (1964); Phys. Rev. **173**, 1452 (1968).

[18] R. Englemann, H. Filthuth, V. Hepp, and E. Kluge, Phys. Lett. **21**, 587 (1966).

[19] B. Sechi-Zorn, B. Kehoe, J. Twitty, and R. A. Barnstein, Phys. Rev. **175**, 1735 (1968).

[20] J. A. Kadyk, G. Alexander, J. H. Chan, P. Gaposchkin, and G. H. Trilling, Nucl. Phys. **B27**, 13 (1971).

[21] M. M. Nagels, T. A. Rijken, and J. J. de Swart, Phys. Rev. D **15**, 2547 (1977).

[22] M. M. Nagels, T. A. Rijken, and J. J, de Swart, Phys. Rev. D **20**, 1633 (1979).

[23] J. J. de Swart, Proceedings of the *U. S. – Japan Seminar on the Hyperon-Nucleon Interaction*, ed. by B. F. Gibson, P. D. Barnes, and K. Nakai, (World Scientific, Singapore, 1994), pp.

[24] R. Timmermans, Proceedings of the *U. S. – Japan Seminar on the Hyperon-Nucleon Interaction*, ed. by B. F. Gibson, P. D. Barnes, and K. Nakai, (World Scientific, Singapore, 1994), pp.

[25] B. Holzenkamp, K. Holinde, and J. Speth, Nucl. Phys. **A500**, 485 (1989).

[26] K. Holinde, Nuc. Phys. **A547**, 255c (1992).

[27] A. G. Reuber, K. Holinde, and J. Speth, Czech. J. Phys. **42**, 1115 (1992).

[28] A. G. Reuber, Proceedings of the *U. S. – Japan Seminar on the Hyperon-Nucleon Interaction*, ed. by B. F. Gibson, P. D. Barnes, and K. Nakai, (World Scientific, Singapore, 1994), pp.

[29] M. Oka, Proceedings of the *U. S. – Japan Seminar on the Hyperon-Nucleon Interaction*, ed. by B. F. Gibson, P. D. Barnes, and K. Nakai, (World Scientific, Singapore, 1994), pp.

[30] B. F. Gibson and D. R. Lehman, Nucl. Phys. **A329**, 308 (1979).

[31] B. F. Gibson and G. J. Stephenson, Phys. Rev. C **8**, 1222 (1973); **11**, 1448 (1975).

[32] B. F. Gibson and D. R. Lehman, Phys. Rev. C **14**, 685 (1976); **15**, 2257 (1977).

[33] L. H. Thomas, Phys. Rev. **47**, 903 (1935); H. A. Bethe and R. F. Bacher, Rev. Mod. Phys. **8**, 82 (1936).

[34] W. Stepien-Rudzka and S. Wycech, Nucl. Phys. A **362**, 349 (1981).

[35] B. F. Gibson and D. R. Lehman, Phys. Rev. C **37** 679 (1988).

[36] R. E. Chrien, S. Bart, M. May, P. H. Pile, R. J. Sutter, P. Barnes, B. Bassalleck, R. Eisenstein, G. Franklin, R. Grace, D. Marlow, R. Rieder, *et al.* Phys. Rev. C **41**, 1062 (1990).

[37] R. H. Dalitz and A. Gal, Ann. Phys. (N.Y.) **116**, 167 (1978).

[38] D. J. Millener, A. Gal, C. B. Dover, and R. H. Dalitz, Phys. Rev. C **31**, 499 (1985).

[39] J. J. de Swart and C. Iddings, Phys. Rev. **128**, 2810 (1962).

[40] L. H. Schick and A. J. Toepfer, Phys. Rev. **170**, 946 (1968); **175**, 1253 (1968).

[41] K. Miyagawa and W. Glöckle, Phys. Rev. C **48**, 2576 (1993).

[42] B. F. Gibson, A. Goldberg, and M. S. Weiss, Phys. Rev. C **6**, 741 (1972).

[43] B. F. Gibson, A. Goldberg, and M. S. Weiss, in *Few Particle Problems in Nuclear Interactions*, (North Holland, Amsterdam, 1972), p. 188.

[44] M. M. Nagels, T. A. Rijken, and J. J. de Swart, Phys. Rev. D **17**, 768 (1978).

[45] J. A. Carlson, V. R. Pandharipande, and R. B. Wiringa, Nucl. Phys. **A401**, 59 (1983).

[46] D. J. Prowse, Phys. Rev. Lett. **17**, 782 (1966).

[47] R. H. Dalitz, D. H. Davis, P. H. Fowler, A. Montwill, J. Pniewski, and J. A. Zakrewski, Proc. Roy. Soc. London **A426**, 1 (1989).

[48] B. F. Gibson, A. Goldberg, and M. S. Weiss, Phys. Rev. **181**, 1486 (1969).

[49] A. R. Bodmer, Q. N. Usmani, and J. A. Carlson, Nucl. Phys. **A422**, 510 (1984).

[50] H. Bandō, Prog. Theor. Phys. **67**, 669 (1982).

[51] M. Danysz, K. Garbowska, J. Pniewski, T. Pniewski, J. Zakrewski, E. R. Fletcher, *et al.*, Phys. Rev. Lett. **11**, 29 (1963); Nucl. Phys. **49**, 121 (1963).

[52] S. Aoki, S. Y. Bahk, K. S. Chung, S. H. Chung, H. Funahashi, C. Hahn, *et al.* Prog. Theo. Phys. **85**, 1287 (1991).

[53] C. B. Dover, D. J. Millener, A. Gal, and D. H. Davis, Phys. Rev. C **44**, 1905 (1991).

This figure "fig1-1.png" is available in "png" format from:

http://arXiv.org/ps/nucl-th/9411027v1



(a)



(b)



(c)